# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



# INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

	11) International Publication Number: WO 88/ 08704
A2	43) International Publication Date: 17 November 1988 (17.11.88)
•	boratories Limited, Mundells, Welwyn Garden City,
87109	(81) Designated States: AU, DK, JP, KR, US.
•	Published
IMIT	· ·
ane, D ONAR oad, M .N (GB	
	1   `

(54) Title: PHARMACEUTICAL COMPOSITIONS

#### (57) Abstract

The invention provides a solid pharmaceutical dosage form comprising: (i) cimetidine; and (ii) antacid, wherein at least part of the antacid is in the form of granules comprising a freely water-soluble solid diluent, the antacid, and a rapidly swellable water-insoluble disintegrant. Compositions of this type overcome the problem of the reduced bioavailability of cimetidine which can occur when cimetidine is co-administered with antacids.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA.	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria.	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil ·	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic		of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	
CH	Switzerland	LI	Liechtenstein	SU	Senegal
CM	Cameroon	LK	Sri Lanka		Soviet Union
DE	Germany, Federal Republic of			TD	Chad
		LU	Luxembourg	TG	Togo
DK	Denmark	MÇ	Monaco	US	United States of America
FI	Finland	MG	Madagascar	•	

10

#### PHARMACEUTICAL COMPOSITIONS

This invention relates to a solid pharmaceutical dosage form comprising cimetidine and an antacid and to a method for the preparation of such a dosage form.

Cimetidine is a histamine H<sub>2</sub>-antagonist which has been described in U.K. Patent Specification 1,397,436. Cimetidine has been shown to be useful in the treatment of duodenal, gastric, recurrent and stomal ulceration, and reflux oesophagitis and in the management of patients who are at high risk from haemorrhage of the upper gastrointestinal tract.

Cimetidine and antacids are frequently
co-administered (see for example the article by
H. Allgayer and G. Paumgartner, Arzneim Forsch.
pp.870-871, 34, No. 8 (1984)). The rationale for
co-administration is that antacid brings about rapid
relief from the symptoms of excess stomach acidity by
neutralising the acid whereas the cimetidine brings about
more sustained relief by inhibiting secretion of more
acid.

25 However, it is well known (see Allgayer and Paumgartner, and Steinberg et al, New England J. Medicine, 1982; 307, 400-4) that when cimetidine is co-administered with antacids, particularly aluminium hydroxide and magnesium hydroxide, there is frequently a 30 substantial reduction in the bioavailability of the The reason for the reduction in cimetidine. bioavailability is not clear, although a number of attempts to discover the mechanism responsible for the problem have been reported in the literature. 35 example Allgayer and Paumgartner were unable to demonstrate why the decrease in bioavailability occurs

25

although they indicated that it was not due to binding of the cimetidine by the antacid.

The benefits, particularly in terms of patient compliance with a treatment regimen, which would arise from an effective combination product containing cimetidine and an antacid, would be expected to be considerable. However, the problem of loss of bioavailability, and the lack of understanding of its cause have, up until now, precluded the development of such a product, as far as we are aware.

It has now been found, surprisingly, that the problem of the loss of bioavailability of the cimetidine can be solved by granulating at least part of the antacid separately, and in a particular manner, prior to mixing with the cimetidine.

In a first aspect, therefore, the invention provides a solid pharmaceutical dosage form comprising:

- (i) cimetidine: and
- (ii) antacid, wherein at least part of the antacid is in the form of granules comprising a freely water-soluble solid diluent, the antacid, and a rapidly swellable water-insoluble disintegrant.

It is preferred that at least 50% by weight of the total antacid in the dosage form is granulated in the particular manner described above in (ii). In general, as the ratio of antacid to cimetidine is increased, it is

desirable to increase the proportion of antacid granulated in this way. In one preferred embodiment of the invention, substantially all of the antacid is thus granulated.

5

The term "freely soluble" is known in the art as referring to a particular level of solubility; thus, in the U.S. Pharmacopoeia, it is defined as meaning that a substance can form a 10% solution in a solvent.

Preferably the substance can form at least a 50% solution in water.

Typically the freely water-soluble solid diluent is a sugar and/or sugar alcohol.

15

Examples of sugars and sugar alcohols are sucrose, lactose, sorbitol, xylitol and mannitol, preferred diluents being lactose, sorbitol and sorbitol/lactose mixtures.

20

It is preferred that the ratio (w/w) of solid diluent to antacid is in the range 1:1 to 8:1, particularly approximately 3:1.

Typically the rapidly swellable water-insoluble disintegrants are synthetic or semi-synthetic polymers of a type known in the art as superdisintegrants (see for example International Patent Application No. PCT/US 87/00302, published as WO 87/05804, and references cited therein). Examples of superdisintegrants include cross-linked polymeric disintegrants such as cross-linked carboxymethyl celluloses, particularly croscarmellose sodium and croscarmellose calcium, cross-linked polyvinyl pyrrolidone, and sodium starch glycolate.

35

Typically the disintegrant is present in an amount from approximately 0.5% (w/w) to approximately 8% (w/w). relative to the total weight of the granules, particularly approximately 2% (w/w).

5

10

The antacid-containing granules are preferably formed by a dry granulation process, for example by compacting using a roller compacter or a tablet press, followed by milling the compact to give granules that have low friability. Suitably, in such a case, the mixture for granulating can contain a lubricant. Examples of lubricants are stearates such as magnesium stearate, and stearic acid.

The antacid typically is selected from aluminium hydroxide, magnesium hydroxide, magnesium carbonate, calcium carbonate and co-dried gels, for example aluminium hydroxide-magnesium carbonate co-dried gel. A particular antacid is aluminium hydroxide or a mixture of aluminium hydroxide and magnesium hydroxide.

In general a dosage form contains between 5 mEq and 30 mEq of antacid, and preferably approximately 14 mEq.

The cimetidine will usually be present in an amount from 50 mg to 800 mg per dosage form, and typically a dosage form will contain 100 mg or 200 mg of cimetidine.

Examples of dosage forms include tablets, capsules 30 and lozenges.

The compositions of the present invention can be in the form of chewable tablets, that is to say tablets which disintegrate readily in the mouth when chewed. With chewable tablets, the pronounced bitter taste of cimetidine means that in practice it is necessary to

10

15

30

provide a means of masking the bitter taste. One means of masking the bitter taste is to coat the cimetidine with a coating agent in an amount effective to mask the bitter taste but which does not significantly affect the bioavailability of the cimetidine.

One such coating agent is dimethylaminoethylmethacrylate/methacrylic acid ester co-polymer which is
sold under the trade name Eudragit E. According to a
co-pending patent application (reference no. 11940)
claiming priority from British patent applications Nos.
8710965 and 8710966, cimetidine can be granulated using
Eudragit E, in an amount 2 - 20% (w/w) relative to the
cimetidine, as the granulating agent. By employing a
Eudragit E loading in this range, the bitter taste of
cimetidine is masked but the dissolution characteristics
and hence bioavailability remain acceptable.

In addition to cimetidine and antacid-containing
granules, the solid dosage forms of the present invention
can contain other pharmaceutical excipients. Thus, for
example, where the dosage form is subject to a compression
step, the dosage form can additionally contain a
lubricating agent, typically stearic acid or a stearate
salt and particularly magnesium stearate.

The compositions of the present invention can also contain additional sweeteners, for example aspartame, cyclamate and saccharin, and colouring and flavouring agents as known in the art.

The invention will now be illustrated in greater detail by the following Examples.

## 100 mg Chewable Tablet

## Ingredient

<b>5</b>	Cimetidine Premix Granules Cimetidine Eudragit El00*	mg/tablet 100.0 10.0	% % % % % % % % % % % % % % % % % % %
•	Antacid (Al/Mg) Granules	mg/tablet	<u>%W/W</u>
10	Direct Compression Sorbitol	590.0	34.01
	Direct Compression Lactose Crystalline	325.0	18.73 18.73
•	Spray dried .	325.0 30.0	1.73
	Croscarmellose Sodium Type A	250.0	14.41
	Dried Aluminium Hydroxide Gel**	200.0	11.53
	Magnesium Hydroxide** Magnesium Stearate	15.0	0.86
15	Magnesium Stealate	1735.0	100.00
	Tableting Mix for Compression  Cimetidine	mg/tablet	
	Premix Granules	110.0	•
	7-4-5-2 (7.1 /35m)		
20	Antacid (Al/Mg) Granules	1735.0	. •
	Aspartame	3.0	
	Peppermint	15.0	
	Tutti Frutti	5.0	
	Spearmint	5.0	
	Lactose	200.0	-
25	Croscarmellose Sodium Type A	30.0	
	Magnesium Stearate	15.0 2118.0	

<sup>\*</sup> Added to the cimetidine by granulation as a 40% W/V solution in methylene chloride. Solvent lost in processing.

<sup>\*\*</sup> Quantities used adjusted for the potencies of raw materials:
Standard quantity of Dried Aluminium Hydroxide gel is equivalent to 117.5 mg/tablet Al<sub>2</sub>O<sub>3</sub> or 180 mg/tablet Aluminium Hydroxide (Al(OH)<sub>3</sub>).

## Process Description

A 40% w/v solution of the Eudragit Eloo in methylene chloride is added with mixing to the cimetidine and blended until granules are formed. The resulting granules are dried and then sieved through a 16 mesh screen.

The aluminium hydroxide, magnesium hydroxide and other ingredients for the antacid granules are sieved through a 12 mesh (1.4 mm) screen and mixed together. The resulting mix is compressed on a rotary tablet press and the resulting compacts are milled using a 12 mesh screen.

15

20

The cimetidine granules, antacid granules and extragranular excipients are put into a cone blender and mixed thoroughly. The resulting mix is discharged from the blender and compressed on a suitable rotary tablet press fitted with the appropriate punches.

## 200 mg Chewable Tablet

## Ingredient

		•	
5 .	Cimetidine Premix Granules Cimetidine Eudragit El00*	mg/tablet 200.0 20.0	%w/w 90.9 9.1
	Antacid (Al/Mg) Granules		
10		mg/tablet	₹W/W
τά	Sorbitol: Direct Compression Grade Lactose: Direct Compression Grade	295.0	34.01
	Spray dried Crystalline	162.5 162.5	18.73 18.73
	Dried Aluminium Hydroxide Gel	125.0	14.41
	Magnesium Hydroxide Croscarmellose Sodium Type A	100.0 15.0	11.53 1.73
15	Magnesium Stearate	7.5 867.5	0.86 100.00
•	Tableting Mix for Compression	mg/tablet	
20	Cimetidine		
• • •	Premix Granules	220.0	
	Antacid (Al/Mg) Granules	867.5	
	Dried Aluminium Hydroxide Gel Magnesium Hydroxide	125.0 100.0	:
25	Sorbitol: Direct Compression Grade Lactose: Direct Compression Grade	295.0	
	Spray dried Crystalline	162.5 162.5	
	Croscarmellose Sodium Type A Aspartame	45.0 3.0	
3.0	Aniseed Butterscotch	20.0	
30	Magnesium Stearate TOTAL		37.5 2063.0

<sup>\*</sup> Added to the cimetidine by granulation as a 40% w/v solution in methylene chloride. Solvent lost in processing.

35

10

## Process Description

The cimetidine premix granules and antacid granules were prepared according to the method described in Example 1. The cimetidine granules and antacid granules were then blended with the remaining ingredients and compressed on a rotary press fitted with the appropriate tablet punches and dies. The formulations of the following Examples 3 and 4 were prepared in a similar manner.

## 200 mg Chewable Tablet

## Ingredient

5 ··· ·	<u>Cimetidine Premix Granules</u> Cimetidine Eudragit El00*	mg/tablet 200.0 20.0	% w/w 90.9 9.1
	Antacid (Al/Mg) Granules		
10		mg/tablet	₹W/W
	Sorbitol: Direct Compression Grade Lactose: Direct Compression Grade	590.0	34.01
	Spray dried	325.0	18.73
	Crystalline	325.0	18.73
	Dried Aluminium Hydroxide Gel	250.0	14.41
	Magnesium Hydroxide	200.0	11.53
	Croscarmellose Sodium Type A+	30.0	1.73
15 .	Magnesium Stearate	$\frac{15.0}{1735.0}$	$\frac{0.86}{100.00}$
			•
	Tableting Mix for Compression	• .	
•		mg/tablet	
20	Cimetidine Premix Granules	220.0	
,	Antacid (Al/Mg) Granules	1735.0	
•	Microcrystalline Cellulose (Avicel PH102)+	200.0	
25	Aspartame	10.0	•
	Aniseed	20.0	
	Butterscotch	20.0	
	Magnesium Stearate	15.0	
	TOTAL	2220.0	

<sup>30 \*</sup> Added to the cimetidine by granulation as a 40% w/v solution in methylene chloride. Solvent lost in processing.

<sup>+</sup> Croscarmellose Sodium Type A and Avicel PH102 can both be obtained from the FMC Corporation, Philadelphia PA.

## 100 mg Chewable Tablet

## <u>Ingredient</u>

5	<u>Cimetidine Premix Granules</u> Cimetidine Eudragit El00*	mg/tablet 100.0 10.0	%W/W 90.9 9.1
	Antacid (Al/Mg) Granules  Lactose: Direct Compression Grade	mg/tablet	<u>%w/w</u>
10	Spray dried Crystalline Dried Aluminium Hydroxide Gel	190.0 190.0 125.0	29.23 29.23 19.23
	Magnesium Hydroxide Croscarmellose Sodium Type A Magnesium Stearate	100.0 30.0 <u>15.0</u> 650.0	15.38 4.62 2.31 100.00
15	Tableting Mix for Compression	850.0	100.00
	Cimetidine Premix Granules	mg/tablet 110.0	
20	Antacid (Al/Mg) Granules	650.0	
	Dried Aluminium Hydroxide Gel Magnesium Hydroxide Sorbitol: Direct Compression Grade	125.0 100.0 590.0	
	Lactose: Direct Compression Grade Spray dried Crystalline	190.0	
	Croscarmellose Sodium Type A Aspartame Aniseed Butterscotch Magnesium Stearate Sodium Saccharin TOTAL	30.0 3.0 20.0 20.0 15.0 5.0 2048.0	
30	2022		

<sup>\*</sup> Added to the cimetidine by granulation as a 40% w/v solution in methylene chloride. Solvent lost in processing.

#### CLAIMS

- 1. A solid pharmaceutical dosage form comprising:
- (i) cimetidine; and
- (ii) antacid, wherein at least part of the antacid is in the form of granules comprising a freely water-soluble solid diluent, the antacid, and a rapidly swellable water-insoluble disintegrant.
- 2. A solid pharmaceutical dosage form according to claim 1 wherein at least 50% of the total antacid present is contained within the granules.
- 3. A solid pharmaceutical dosage form according to claim 2 wherein substantially all of the antacid present is contained within the granules.
  - 4. A dosage form according to any one of claims 1 to 3 wherein the granules are dry-granulated.

20

- 5. A dosage form according to any one of claims 1 to 4 wherein the highly water-soluble solid diluent is a sugar or a sugar alcohol.
- 6. A dosage form according to any one of claims 1 to 5 wherein the ratio (w/w) of solid diluent to antacid is in the range from approximately 1:1 to approximately 8:1.
- 7. A dosage form according to any one of claims 1 to 6 wherein the disintegrant is a cross-linked carboxymethylcellulose.

ŗ

- 8. A dosage form according to any one of claims 1 to 7 wherein the antacid is present in an amount from between 5 mEq and 30 mEq.
- 9. A chewable pharmaceutical tablet composition comprising:
  - (i) granules comprising cimetidine and a granulating agent wherein the granulating agent is a co-polymer of dimethylaminoethylmethacrylate and methacrylic acid ester in an amount of approximately 10% (w/w) relative to the cimetidine; and
- (ii) antacid-containing granules comprising aluminium hydroxide and magnesium hydroxide; a solid diluent which is lactose or a mixture of sorbitol and lactose, the ratio (w/w) of diluent to aluminium hydroxide/magnesium hydroxide being approximately 3:1; and a disintegrant which is croscarmellose sodium, the disintegrant being present in an amount of approximately 2% (w/w) relative to the total weight of the antacid-containing granules; wherein the antacid-containing granules are formed by dry granulation.
- 10. A granule comprising a highly water-soluble solid diluent, an antacid, and a rapidly swellable water-insoluble disintegrant as defined in any one of claims 1. 4, 5, 7 and 9.

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



#### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)						
(51) International Patent Classification <sup>4</sup> : A61K 31/415, 33/10, 9/20 A61K 33/08	A3 .	(11) International Publication Number: WO 88/08704 (43) International Publication Date: 17 November 1988 (17.11.88)				
(21) International Application Number: PCT/GB (22) International Filing Date: 4 May 1988 (	•	boratories Limited, Mundells, Welwyn Garden City				
(31) Priority Application Number:	87109	65 (81) Designated States: AU, DK, JP, KR, US.				
(32) Priority Date: 8 May 1987 ( (33) Priority Country:  (71) Applicant (for all designated States except US) KLINE & FRENCH LABORATORIES L [GB/GB]; Mundells, Welwyn Garden City, shire AL7 1EY (GB).  (72) Inventors; and (75) Inventors; Applicants (for US only): FRANCE, [GB/GB]; West Lodge, 2 Harmer Green La swell, Hertfordshire AL6 0AD (GB). LEG Graham, Stanley [GB/GB]; 60 Hazelmere Ro shalswick, St. Albans, hertfordshire AL4 9R1 PEARMAIN, Kevin, Edward [GB/GB]; 24 TI Stotfold, Hitchin, Hertfordshire (GB).	: SMITIMITE hertform, Gordine, Dinar ad, Marad, Min (GB)	Published  With international search report.  Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.  (88) Date of publication of the international search report:  9 February 1989 (09.02.89)  on on one of the international search report:  10 June 1989 (09.02.89)				

#### (54) Title: PHARMACEUTICAL COMPOSITIONS

#### (57) Abstract

The invention provides a solid pharmaceutical dosage form comprising: (i) cimetidine; and (ii) antacid, wherein at least part of the antacid is in the form of granules comprising a freely water-soluble solid diluent, the antacid, and a rapidly swellable water-insoluble disintegrant. Compositions of this type overcome the problem of the reduced bioavailability of cimetidine which can occur when cimetidine is co-administered with antacids.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT AU BB BE BG BI BR CF CG CH DE DK FI	Austria Australia Barbados Belgium Bulgaria Benin Brazil Central African Republic Congo Switzerland Cameroon Germany, Federal Republic of Denmark Finland	FR GA GB HU IT JP KP KR LI LK LU MG MG	France Gabon United Kingdom Hungary Italy Japan Democratic People's Republic of Korea Republic of Korea Liechtenstein Sri Lanka Luxembourg Monaco Madagascar	ML MR MW NL NO SD SE SN SU TIG US	Mali Mauritania Malawi Netherlands Norway Romania Sudan Sweden Senegal Soniet Union Chad Togo United States of America
--	---	--	--	---	--

#### INTERNATIONAL SEARCH REPORT

International Application No PCT/GB 88/00350 I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) \* According to International Patent Classification (IPC) or to both National Classification and IPC IPC4: A 61 K 31/415; A 61 K 33/10; A 61 K 9/20; A 61 K 33/08 Minimum Documentation Searched 7 Classification System Classification Symbols IPC4 A 61 K Documentation Searched other than Minimum Documentation to the Extent that such Documents are included in the Fields Searched III. DOCUMENTS CONSIDERED TO BE RELEVANTS Citation of Document, 11 with Indication, where appropriate, of the relevant passages 12 Relevant to Claim No. 13 Category \* EP, A, 0003589 (THE WELLCOME FOUNDATION X 10 LTD) 22 August 1979, see example 1; claims 1-13 Y 1-8 Y EP, A, 0138540 (SMITH KLINE DAUELSBERG) 1-8 24 April 1985, see page 4, lines 18-26; claims 1-10 FR, M, 6885 (SMITH KLINE & FRENCH LAB.) 1-8 Y 21 April 1969, see example 1 Y Chemical Abstracts, vol. 102, 1985, 1-8 (Columbus, Ohio, US) see page 375, abstract no. 137799m & JP, A, 59193825 (SUMITOMO CHEMICAL CO. LTD) 2 November 1984 "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the \* Special categories of cited documents: 10 "A" document defining the general state of the art which is not considered to be of particular relevance invention earlier document but published on or after the international filing date "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such docudocument referring to an oral disclosure, use, exhibition or other means ments, such combination being obvious to a person skilled in the art. document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family IV. CERTIFICATION Date of Mailing of this International Search Report Date of the Actual Completion of the International Search 7 / 01, 89 27th July 1988

Signature of Authorized Officer

P.C.G. VAN DER PUTTEN

EUROPEAN PATENT OFFICE

International Searching Authority

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET						
V. OBSERVATIONS WHERE CERTAI	N CLAIMS WERE FOUND UNSEARCHABLE 1					
	established in respect of certain claims under Article 17(2) (a) for the following reasons:					
this international search report has intractional	late to subject matter not required to be searched by this Authority, namely:					
in the second se						
	•					
	•					
<u></u>	Cate transmission that do not comply with the prescribed requires					
Claim numbers, because they rel	late to parts of the international application that do not comply with the prescribed require- grul international search can be carried out, specifically:					
manta to add the						
	dependent claims and are not drafted in accordance with the second and third sentences of					
Claim numbers, because they are PCT Rule 6.4(a).	dependent claims and are not draited in accordance					
•	or warming to be explice?					
VI. X OBSERVATIONS WHERE UNITY						
This international Searching Authority found a	multiple inventions in this international application as follows:					
1. Claims 1 - 8,10						
2. Claim 9	for further information please see					
Z. CILLIII	form PCT/ISA/206 mailed on 29.08.198					
	• •					
1. X As all required additional search fees we	re timely paid by the applicant, this international search report covers all searchable claims					
of the International application.	search fees were timely paid by the applicant, this international search report covers only					
those claims of the international applicat	ion for which fees were paid, specifically claims:					
3. No required additional search fees were the invention first mentioned in the claim	timely paid by the applicant. Consequently, this international search report is restricted to is; it is covered by claim numbers:					
	to the section of the					
<ol> <li>As all searchable claims could be searche invite payment of any additional fee.</li> </ol>	ed without effort justifying an additional fee, the international Searching Authority did not					
Remark on Protest						
The additional search fees were accompa	anied by applicant's protest.					
X No protest accompanied the payment of						

# ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

GB 8800350

SA 22061

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 11/01/89

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0003589	22-08-79	JP-A- 54119022 AU-A- 4392179 GB-A- 1601833 CA-A- 1117418 AU-B- 525122	14-09-79 16-08-79 04-11-81 02-02-82 21-10-82
EP-A- 0138540	24-04-85	WO-A- 8501656 AU-A- 3503984 JP-T- 61500223 AU-B- 560608 DE-A- 3467703 CA-A- 1239349	25-04-85 07-05-85 06-02-86 09-04-87 07-01-88 19-07-88
FR-M- 6885	·21-04-69	NL-A- 6617127 US-A- 3501571 BE-A- 690679 DE-A- 1617800 FR-A- 1515061 GB-A- 1129260	07-06-67 17-03-70 05-06-67 08-04-71